

P542. (8.7)

①

3. List the ordered pairs in the relations on $\{1, 2, 3\}$ corresponding to these matrices, (where the rows and columns correspond to the integers listed in increasing order).

a)
$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

$(1,1), (1,3), (2,2), (3,1), (3,3)$

b)
$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

$(1,2), (2,2), (3,2)$

c)
$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

$(1,1), (1,2), (1,3),$

$(2,1), (2,3),$

$(3,1), (3,2), (3,3)$

P543.

1) B. Let R be the relation represented by the matrix

$$M_R = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

Find the matrix representing:

a) R^{-1}

sol: $\begin{bmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$

b) \bar{R}

sol: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$

c) R^2

sol: $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$

P553 (8.4)

3. Let R be the relation $\{(a, b) \mid a \text{ divides } b\}$ on the set of integers. What is the symmetric closure of R ?

solution: $R \cup R^{-1}$
 $= \{(a, b) \mid a \text{ divides } b \text{ or } b \text{ divides } a\}$

P527.

81:3: Let $S = \{1, 2, 3, 4\}$. Determine whether each relation R on S has the listed properties.

$$R = \{(1,1), (1,2), (2,1), (2,2), (3,3), (4,4)\}$$


~~Solution~~
Question 1: Draw the 0-1 matrix representation of R .

Solution:

$$\begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

2: Draw the digraph representation of R .

Solution:



3. Is R symmetric?

Yes

4. Is R transitive?

Yes.

5. Is R reflexive?

Yes

P29. f3.

a) How many relations are there on the set $\{a, b, c, d\}$

$$2^n = 2^{4^2} = 2^{16} = 65536$$

(8.5)

P563 9. Suppose that A is a nonempty set, and f is a function that has A as its domain. Let R be the relation on A consisting of all ordered pairs (x, y) such that $f(x) = f(y)$.

a) Show that R is an equivalence relation on A .

Solution: $(x, x) \in R$ because $f(x) = f(x)$.

Hence, R is reflexive.

$(x, y) \in R$ iff $f(x) = f(y)$, which holds iff $f(y) = f(x)$ iff $(y, x) \in R$.

Hence, R is symmetric.

If $(x, y) \in R$, and $(y, z) \in R$, then $f(x) = f(y)$ and $f(y) = f(z)$. Hence, $f(x) = f(z)$. Thus, $(x, z) \in R$.

It follows that R is transitive.

b) What are the equivalence classes of R ?

the sets $f^{-1}(b)$ for b in the range of f .

P563

8.5 : 3 Which of these relations on the set of all functions from \mathbb{Z} to \mathbb{Z} are equivalence relations? Determine the properties of an equivalence relation that the others lack

a) $\{ (f, g) \mid f(i) = g(i) \}$

Solution \rightarrow It is ~~reflexive~~ transitive.

\because if we have $f R g$, i.e. $f(i) = g(i)$

and $g R h$, i.e. $g(i) = h(i)$

then we will have $f(i) = g(i) = h(i) \Rightarrow f(i) = h(i)$
then we have $f R h$.

\rightarrow It is Reflexive.

\because if we have $f(i) = f(i)$, which implies $f R f$.
thus, it is reflexive.

\rightarrow It is symmetric:

\because if we have $f R g$, i.e. $f(i) = g(i)$

then we will have $g(i) = f(i)$, which implies
 $g R f$.

So, if $f R g$, ~~then~~ then $g R f$. \therefore it is symmetric.

\rightarrow Since R has all three properties,
it is an equivalence relation.

(b) $\{(f, g) \mid f(0) = g(0) \text{ or } f(1) = g(1)\}$.

Solution: \rightarrow It is reflexive:

\because we have $f(0) = f(0)$ or $f(1) = f(1)$ then $f R f$.
so. it is reflexive.

\rightarrow It is symmetric.

\because if we have $f R g$, i.e. $f(0) = g(0)$ or $f(1) = g(1)$
then we have $g(0) = f(0)$ or $g(1) = f(1)$.
thus. $g R f$. so it is symmetric.

\rightarrow It is not transitive.

\because if we have. $f R g$ i.e. $f(0) = g(0)$ or $f(1) = g(1)$
and $g R h$ i.e. $g(0) = h(0)$ or $g(1) = h(1)$

Counter example is,

when $f(0) = g(0) \wedge f(1) \neq g(1)$ i.e. $f R g$
 $g(0) \neq h(0) \wedge g(1) = h(1)$ i.e. $g R h$.

then. we have $f(0) \neq h(0) \wedge f(1) \neq h(1)$
then. $f \not R h$.

\therefore it is not transitive.

\rightarrow Since R does not have all three properties.
It is not an equivalence relation.

c) $\{ (f, g) \mid f(x) - g(x) = 1 \text{ for all } x \in \mathbb{Z} \}$

solution: \rightarrow It is not reflexive.

\therefore if we have $f R f$, then, we should have
 $f(x) - f(x) = 1$ which is impossible.

So, R is not reflexive, which ~~means~~ means $f \not R f$.

\rightarrow It is not symmetric.

\therefore If we have $f R g$, i.e. $f(x) - g(x) = 1$
then, $g(x) - f(x) = -1$, which means $g \not R f$
 \therefore it is not symmetric.

\rightarrow It is not transitive.

\therefore If we have $f R g$, i.e. $f(x) - g(x) = 1$
and we have $g R h$, i.e. $g(x) - h(x) = 1$

$$\therefore f(x) - g(x) + g(x) - h(x) = 1 + 1$$

$$\Rightarrow f(x) - h(x) = 2$$

$$\Rightarrow f \not R h$$

\therefore it is not transitive.

\rightarrow Since R does not have all three properties,
then R is not an equivalence relation.

P538 33. (8.1)

* For the following relations on the set of real numbers:

$R_1 = \{(a, b) \in \mathbb{R}^2 \mid a > b\}$, the "greater than" relation,

$R_2 = \{(a, b) \in \mathbb{R}^2 \mid a \geq b\}$, the "greater than or equal to" relation,

$R_3 = \{(a, b) \in \mathbb{R}^2 \mid a < b\}$, the "less than" relation,

$R_4 = \{(a, b) \in \mathbb{R}^2 \mid a \leq b\}$, the "less than or equal to" relation,

$R_5 = \{(a, b) \in \mathbb{R}^2 \mid a = b\}$, the "equal to" relation,

$R_6 = \{(a, b) \in \mathbb{R}^2 \mid a \neq b\}$, the "unequal to" relation.

Find:

a) $R_2 \cup R_4$
 $= \mathbb{R}^2$

b) $R_3 \cup R_6$
 $= R_6$ ($\because R_6$ includes all relations in R_3)

c) $R_3 \cap R_6$
 $= R_3$

d) $R_4 \cap R_6$
 $= \{(a, b) \in \mathbb{R}^2 \mid a \leq b\} \cap \{(a, b) \in \mathbb{R}^2 \mid a \neq b\}$
 $= \{(a, b) \in \mathbb{R}^2 \mid a < b\}$
 $= R_3$

e) $R_3 - R_6$
 $= \{(a, b) \in \mathbb{R}^2 \mid a < b\} - \{(a, b) \in \mathbb{R}^2 \mid a \neq b\}$
 $= \emptyset$

$$(f) R_6 - R_3$$

$$= \{(a,b) \in \mathbb{R}^2 \mid a \neq b\} - \{(a,b) \in \mathbb{R}^2 \mid a < b\}$$

$$= \{(a,b) \in \mathbb{R}^2 \mid a > b\}$$

$$= R_1$$

$$(g) R_2 \oplus R_6$$

$$= \{(a,b) \in \mathbb{R}^2 \mid a \geq b\} \oplus \{(a,b) \in \mathbb{R}^2 \mid a \neq b\}$$

$$= \{(a,b) \in \mathbb{R}^2 \mid a \leq b\}$$

$$= R_4$$

$$R_2 \oplus R_6 \\ = R_2 \cup R_6 - R_2 \cap R_6$$

$$(h) R_3 \oplus R_5$$

$$= \{(a,b) \in \mathbb{R}^2 \mid a < b\} \oplus \{(a,b) \in \mathbb{R}^2 \mid a = b\}$$

$$= \{(a,b) \in \mathbb{R}^2 \mid a \leq b\}$$

$$= R_4$$

$$R_3 \oplus R_5 \\ = R_3 \cup R_5 - R_3 \cap R_5$$